

Durham Research Online

Deposited in DRO:

06 January 2015

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Belmonte, R. and Hof, van 't P. and Kaminski, M. and Paulusma, D. and Thilikos, D.M. (2013)
'Characterizing graphs of small carving-width.', Discrete applied mathematics., 161 (13-14). pp. 1888-1893.

Further information on publisher's website:

<http://dx.doi.org/10.1016/j.dam.2013.02.036>

Publisher's copyright statement:

NOTICE: this is the author's version of a work that was accepted for publication in Discrete applied mathematics. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Discrete applied mathematics, 161, 13-14, 2013, 10.1016/j.dam.2013.02.036

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Characterizing Graphs of Small Carving-Width*

Rémy Belmonte^{1,†}, Pim van 't Hof^{1,†}, Marcin Kamiński²,
Daniël Paulusma^{3,‡}, and Dimitrios M. Thilikos^{4,§}

¹ Department of Informatics, University of Bergen, Norway
{remy.belmonte,pim.vanthof}@ii.uib.no

² Département d'Informatique, Université Libre de Bruxelles, Belgium
Institute of Computer Science, University of Warsaw, Poland
mjk@mimuw.edu.pl

³ School of Engineering and Computing Sciences, Durham University, UK
daniel.paulusma@durham.ac.uk

⁴ Department of Mathematics, National & Kapodistrian University of Athens, Greece
sedthilk@math.uoa.gr

Abstract. We characterize all graphs that have carving-width at most k for $k = 1, 2, 3$. In particular, we show that a graph has carving-width at most 3 if and only if it has maximum degree at most 3 and treewidth at most 2. This enables us to identify the immersion obstruction set for graphs of carving-width at most 3.

1 Introduction

All graphs considered in this paper are finite and undirected, have no self-loops but may have multiple edges. A graph that has no multiple edges is called *simple*. For undefined graph terminology we refer to the textbook of Diestel [7]. A *carving* of a graph G is a tree T whose internal vertices all have degree 3 and whose leaves correspond to the vertices of G . For every edge e of T , deleting e from T yields exactly two trees, whose leaves define a bipartition of the vertices of G ; we say that the edge cut in G corresponding to this bipartition is *induced by e* . The *width* of a carving T is the maximum size of an edge cut in G that is induced by an edge of T . The *carving-width* of G is the minimum width of a carving of G .

Carving-width was introduced by Seymour and Thomas [17], who proved that checking whether the carving-width of a graph is at most k is an NP-complete problem. In the same paper, they proved that there is a polynomial-time algorithm for computing the carving-width of planar graphs. Later, the problem of constructing carvings of minimum width was studied by Khuller [12],

*The results of this paper have appeared in the proceedings of COCOA 2012 [3].

[†]Supported by the Research Council of Norway (197548/F20).

[‡]Supported by EPSRC (EP/G043434/1) and Royal Society (JP100692).

[§]Co-financed by the European Union (European Social Fund - ESF) and Greek national funds through the Operational Program “Education and Lifelong Learning” of the National Strategic Reference Framework (NSRF) - Research Funding Program: “Thales. Investing in knowledge society through the European Social Fund.”

who presented a polynomial-time algorithm for constructing a carving T whose width is within a $O(\log n)$ factor from the optimal. In [20] an algorithm was given that decides, in $f(k) \cdot n$ steps, whether an n -vertex graph G has carving-width at most k and, if so, also outputs a corresponding carving of G . We stress that the values of $f(k)$ in the complexity of the algorithm in [20] are huge, which makes the algorithm highly impractical even for trivial values of k .

A graph G contains a graph H as an immersion if H can be obtained from some subgraph of G after lifting a number of edges (see Section 2 for the complete definition). Recently, the immersion relation attracted a lot of attention both from the combinatorial [1,6,9,21] and the algorithmic [10,11] point of view. It can easily be observed (cf. [20]) that carving-width is a parameter closed under taking immersions, i.e., the carving-width of a graph is not smaller than the carving-width of any of its immersions. Combining this fact with the seminal result of Robertson and Seymour in [15] stating that graphs are well-quasi-ordered with respect to the immersion relation, it follows that the set \mathcal{G}_k of graphs with carving-width at most k can be completely characterized by forbidding a finite set of graphs as immersions. This set is called an *immersion obstruction set* for the class \mathcal{G}_k .

Identifying obstruction sets is a classic problem in structural graph theory, and its difficulty may vary, depending on the considered graph class. While obstructions have been extensively studied for parameters that are closed under minors (see [2,5,8,13,14,16,18,19] for a sample of such results), no obstruction characterization is known for any immersion-closed graph class. In this paper, we make a first step in this direction.

The outcome of our results is the identification of the immersion obstruction set for \mathcal{G}_k when $k \leq 3$; the obstruction set for the non-trivial case $k = 3$ is depicted in Figure 3. Our proof for this case is based on a combinatorial result stating that \mathcal{G}_3 consists of exactly the graphs with maximum degree at most 3 and treewidth at most 2. A direct implication of our results is a linear-time algorithm for the recognition of the class \mathcal{G}_k when $k = 1, 2, 3$. This can be seen as a “tailor-made” alternative to the general algorithm of [20] for elementary values of k .

2 Preliminaries

Let $G = (V, E)$ be a graph, and let $S \subset V$ be a subset of vertices of G . Then the set of edges between S and $V \setminus S$, denoted by $(S, V \setminus S)$, is an *edge cut* of G . Let the vertices of G be in 1-to-1 correspondence to the leaves of a tree T whose internal vertices all have degree 3. The correspondence between the leaves of T and the vertices of G uniquely defines the following edge weighting w on the edges of T . Let $e \in E_T$, and let C_1 and C_2 be the two connected components of $T - e$. Let S_i be the set of leaves of T that are in C_i for $i = 1, 2$; note that $S_2 = V \setminus S_1$. Then the weight $w(e)$ of the edge e in T is the number of edges in the edge cut (S_1, S_2) of G . The tree T is called a *carving* of G , and (T, w) is a *carving decomposition* of G . The *width* of a carving decomposition (T, w) is

the maximum weight $w(e)$ over all $e \in E_T$. The *carving-width* of G , denoted by $\text{cw}(G)$, is the minimum width over all carving decompositions of G . We define $\text{cw}(G) = 0$ if $|V| = 1$. We refer to Figure 4 for an example of a graph and a carving decomposition.

A *tree decomposition* of a graph $G = (V, E)$ is a pair $(\mathcal{T}, \mathcal{X})$, where \mathcal{X} is a collection of subsets of V , called *bags*, and \mathcal{T} is a tree whose vertices, called *nodes*, are the sets of \mathcal{X} , such that the following three properties are satisfied:

- (i) for each $u \in V$, there is a bag $X \in \mathcal{X}$ with $u \in X$;
- (ii) for each $uv \in E$, there is a bag $X \in \mathcal{X}$ with $u, v \in X$;
- (iii) for each $u \in V$, the nodes containing u induce a connected subtree of \mathcal{T} .

The *width* of a tree decomposition $(\mathcal{T}, \mathcal{X})$ is the size of a largest bag in \mathcal{X} minus 1. The *treewidth* of G , denoted by $\text{tw}(G)$, is the minimum width over all possible tree decompositions of G .

Let $G = (V, E)$ be a graph, and let uv be an edge of G . The *contraction* of the edge uv is the operation that deletes u and v from G and replaces them by a new vertex x that is made adjacent to the neighbors of u and of v in G , such that for every vertex $w \in V \setminus \{u, v\}$, the number of edges between x and w in the new graph is equal to the number of edges between w and $\{u, v\}$ in G . A graph G contains a graph H as a *minor* if H can be obtained from G by a sequence of vertex deletions, edge deletions and edge contractions. The following two well-known properties of treewidth will be used in the proof of our main result.

Lemma 1 (cf. [4]). *Let G be a simple graph and k an integer. If $\text{tw}(G) \leq k$, then G contains a vertex of degree at most k .*

Lemma 2 (cf. [4,7]). *Let G be a graph. Then $\text{tw}(H) \leq \text{tw}(G)$ for every minor H of G .*

The *subdivision* of an edge uv is the operation that deletes the edge uv from the graph and adds a new vertex w as well as two new edges uw and vw . The reverse operation is called *vertex dissolution*; this operation removes a vertex v of degree 2 that has two distinct neighbors u and w , and adds a new edge between u and w , regardless of whether or not there already exist edges between u and w . A graph G contains a graph H as a *topological minor* if H can be obtained from G by a sequence of vertex deletions, edge deletions, and vertex dissolutions. Equivalently, G contains H as a topological minor if G contains a subgraph H' that is a *subdivision* of H , i.e., H' can be obtained from H by a sequence of edge subdivisions. The following lemma is obtained by combining some well-known properties of treewidth, minors, and topological minors.

Lemma 3 (cf. [7]). *A graph has treewidth at most 2 if and only if it does not contain K_4 as a topological minor.*

Let u, v, w be three distinct vertices in a graph such that uv and vw are edges. The operation that removes the edges uv and vw , and adds the edge uw (even

in the case u and w are already adjacent) is called a *lift*. A graph G contains a graph H as an *immersion* if H can be obtained from G by a sequence of vertex deletions, edge deletions, and lifts. Note that dissolving a vertex v of degree 2 that has two distinct neighbors u and w is equivalent to first lifting the edges uv and vw and then deleting the vertex v . Hence, it readily follows from the definitions of topological minors and immersions that every topological minor of graph G is also an immersion of G .

3 The Main Result

We begin this section by stating some useful properties of carving-width. The following observation is known and easy to verify by considering the number of edges in the edge cut $(\{u\}, V \setminus \{u\})$ of a graph $G = (V, E)$.

Observation 1 *Let G be a graph. Then $\text{cw}(G) \geq \Delta(G)$.*

We also need the following two straightforward lemmas. The first lemma follows from the observation that any subgraph of a graph is an immersion of that graph, combined with the observation that carving-width is a parameter that is closed under taking immersions (cf. [20]). We include the proof of the second lemma for completeness.

Lemma 4. *Let G be a graph. Then $\text{cw}(H) \leq \text{cw}(G)$ for every subgraph H of G .*

Lemma 5. *Let G be a graph with connected components G_1, \dots, G_p for some integer $p \geq 1$. Then $\text{cw}(G) = \max\{\text{cw}(G_i) \mid 1 \leq i \leq p\}$.*

Proof. Since the carving-width of a graph with only one vertex is defined to be 0, the lemma clearly holds if G has no edges. Suppose G has at least one edge. Lemma 4 implies that $\max\{\text{cw}(G_i) \mid 1 \leq i \leq p\} \leq \text{cw}(G)$. Now let (T_i, w_i) be a carving decomposition of G_i of width $\text{cw}(G_i)$ for $i = 1, \dots, p$. Since deleting isolated vertices does not change the carving-width of a graph, we may without loss of generality assume that G has no isolated vertices. In particular, this means that each tree T_i contains at least one edge. We construct a carving decomposition (T, w) of G from the p carving decompositions (T_i, w_i) as follows.

We pick an arbitrary edge $e_i = x_i y_i$ in each T_i . For each $i \in \{2, \dots, p-1\}$, we subdivide the edge e_i twice by replacing it with edges $x_i z_i$, $z_i z'_i$ and $z'_i y_i$, where z_i and z'_i are two new vertices. The edges e_1 and e_p are subdivided only once: the edge e_1 is replaced with a new vertex z_1 and two new edges $x_1 z_1$ and $z_1 y_1$, and the edge e_p is replaced with a new vertex z'_p and two new edges $x_p z'_p$ and $z'_p y_p$. Finally, we add the edge $z_i z'_{i+1}$ for each $i \in \{1, \dots, p-1\}$. This results in a tree T whose internal vertices all have degree 3. Since there are no edges between any two connected components of G , the corresponding carving decomposition (T, w) of G has width $\max\{\text{cw}(G_i) \mid 1 \leq i \leq p\}$. Hence, $\text{cw}(G) \leq \max\{\text{cw}(G_i) \mid 1 \leq i \leq p\}$. We conclude that $\text{cw}(G) = \max\{\text{cw}(G_i) \mid 1 \leq i \leq p\}$. \square

The next lemma is the final lemma we need in order to prove our main result.




Fig. 1. A schematic illustration of how the tree T' in the carving decomposition of G' is transformed into a tree T in the proof of Lemma 6 when the edge uw in G' is subdivided. The vertex x is an arbitrary vertex of G' , possibly w .

Lemma 6. *Let G' be a graph with carving-width at least 2, and let uw be an edge of G' . Let G be the graph obtained from G' by subdividing the edge uw . Then $\text{cw}(G) = \text{cw}(G')$.*

Proof. Let (T', w') be a carving decomposition of G' of width $\text{cw}(G') \geq 2$, and let p be the unique neighbor of u in T' . Let v be the vertex that was used to subdivide the edge uw in G' , i.e., the graph G was obtained from G' by replacing uw with edges uv and vw for some new vertex v . Let T be the tree obtained from T' by first relabeling the leaf in T' corresponding to vertex u by q , and then adding two new vertices u and v as well as two new edges qu and qv ; see Figure 1 for an illustration. Let us show that the resulting carving decomposition (T, w) of G has width at most $\text{cw}(G')$.

Let e be an edge in T . Suppose that $e = pq$. By definition, $w(e)$ is the number of edges between $\{u, v\}$ and $V \setminus \{u, v\}$ in G , which is equal to the number of edges incident with u in G' . The latter number is the weight of the edge pu in T' . Hence, $w(e) \leq \text{cw}(G')$. Suppose that $e = qu$. By definition, $w(e)$ is the number of edges incident with u in G , which is equal to the number of edges incident with u in G' . Hence $w(e) \leq \text{cw}(G')$. Suppose that $e = qv$. By definition, $w(e)$ is the number of edges incident with v in G , which is 2. Hence $w(e) = 2 \leq \text{cw}(G')$. Finally, suppose that $e \notin \{pq, qu, qv\}$. Let C_1 and C_2 denote the subtrees of T obtained after removing e . Let S_i be the set of leaves of T in C_i for $i = 1, 2$. Then u and v either both belong to S_1 or both belong to S_2 . Without loss of generality, assume that both u and v belong to S_1 . By definition, $w(e)$ is the number of edges between S_1 and S_2 in G , which is equal to the number of edges between $S_1 \setminus \{v\}$ and S_2 in G' . The latter number is the weight of the edge e in

T' . Hence, $w(e) \leq \text{cw}(G')$. We conclude that (T, w) has width at most $\text{cw}(G')$, and hence $\text{cw}(G) \leq \text{cw}(G')$.

It remains to show that $\text{cw}(G) \geq \text{cw}(G')$. Let (T^*, w^*) be a carving decomposition of G of width $\text{cw}(G)$. We remove the leaf corresponding to v from T^* . Afterwards, the neighbor of v in T^* has degree 2, and we dissolve this vertex. This results in a tree T'' . It is easy to see that the corresponding carving decomposition (T'', w'') of G' has width at most $\text{cw}(G)$. Hence, $\text{cw}(G) \geq \text{cw}(G')$. This completes the proof of Lemma 6. \square

We are now ready to show the main result of our paper.

Theorem 1. *Let G be a graph. Then the following three statements hold.*

- (i) $\text{cw}(G) \leq 1$ if and only if $\Delta(G) \leq 1$.
- (ii) $\text{cw}(G) \leq 2$ if and only if $\Delta(G) \leq 2$.
- (iii) $\text{cw}(G) \leq 3$ if and only if $\Delta(G) \leq 3$ and $\text{tw}(G) \leq 2$.

Proof. Let $G = (V, E)$ be a graph. By Lemma 5 we may assume that G is connected. We prove the three statements separately.

(i) If $\text{cw}(G) \leq 1$, then $\Delta(G) \leq 1$ due to Observation 1. If $\Delta(G) \leq 1$, then G is isomorphic to either K_1 or K_2 . Clearly, $\text{cw}(G) \leq 1$ in both cases.

(ii) If $\text{cw}(G) \leq 2$, then $\Delta(G) \leq 2$ due to Observation 1. If $\Delta(G) = 1$, then $\text{cw}(G) \leq 1$ follows from (i). If $\Delta(G) = 2$, then G is either a graph consisting of two vertices with two edges between them, or a simple graph that is either a path or a cycle. In all three cases, it is clear that $\text{cw}(G) \leq 2$.

(iii) First suppose that $\text{cw}(G) \leq 3$. Then $\Delta(G) \leq 3$ due to Observation 1. We need to show that $\text{tw}(G) \leq 2$. For contradiction, suppose that $\text{tw}(G) \geq 3$. Then, by Lemma 3, G contains K_4 as a topological minor, i.e., G contains a subgraph H such that H is a subdivision of K_4 . Since $\text{cw}(K_4) = 4$, we have that $\text{cw}(H) = \text{cw}(K_4) = 4$ as a result of Lemma 6. Since H is a subgraph of G , Lemma 4 implies that $\text{cw}(G) \geq \text{cw}(H) = 4$, contradicting the assumption that $\text{cw}(G) \leq 3$.

For the reverse direction, we need to prove that every graph $G = (V, E)$ with $\Delta(G) \leq 3$ and $\text{tw}(G) \leq 2$ has carving-width at most 3. We use induction on $|V|$. If $|V| \leq 2$, then G is either isomorphic to K_1 or K_2 , or G consists of two vertices with exactly two edges between them. It is clear that $\text{cw}(G) \leq 3$ in each of these cases. From now on, we assume that $|V| \geq 3$.

First, suppose that G contains two vertices u and v with at least two edges between them. Since $|V| \geq 3$, we may without loss of generality assume that v has a neighbor $t \neq u$. Then, because $\Delta(G) \leq 3$ and there are at least two edges between u and v in G , we find that t and u are the only two neighbors of v in G and that the number of edges between u and v is exactly 2. Let G^* denote the graph obtained from G by deleting one edge between u and v , and let G' denote the graph obtained from G^* by dissolving v . Note that G' is a connected graph on $|V| - 1$ vertices, and $\Delta(G') \leq 3$. Moreover, since G' is a topological minor and

hence a minor of G , Lemma 2 ensures that $\text{tw}(G') \leq \text{tw}(G) \leq 2$. Consequently, we can apply the induction hypothesis to deduce that $\text{cw}(G') \leq 3$.

If $\text{cw}(G') \leq 1$, then $\Delta(G') \leq 1$ by Observation 1. Since G' is a connected graph on $|V| - 1 \geq 2$ vertices, G' must be a path on two vertices. Then G^* is a path on three vertices, implying that $\text{cw}(G^*) = 2$. Since G can be obtained from G^* by adding a single edge, $\text{cw}(G) \leq 3$ in this case. Suppose $2 \leq \text{cw}(G') \leq 3$. Then, by Lemma 6, $\text{cw}(G^*) = \text{cw}(G') \leq 3$. Moreover, from the proof of Lemma 6 it is clear that there exists a carving decomposition (T^*, w^*) of G^* of width $\text{cw}(G^*)$ such that u and v have a common neighbor q in T^* . We consider the carving decomposition (T, w) of G with $T = T^*$. Let e be an edge in T . First suppose that $e = uq$ or $e = vq$. Then $w(e) \leq 3$, as both u and v have degree at most 3 in G . Now suppose that $e \notin \{uq, vq\}$. Then $w(e) = w^*(e) \leq \text{cw}(G^*) \leq 3$. We conclude that the carving decomposition (T, w) of G has width at most 3, which implies that $\text{cw}(G) \leq 3$.

From now on, we assume that G contains no multiple edges, i.e., we assume that G is simple. Since $\text{tw}(G) \leq 2$, G contains a vertex of degree at most 2 due to Lemma 1.

Suppose G contains a vertex u of degree 1. Let v be the neighbor of u in G , and let G' be the graph obtained from G by deleting u . It is clear that $\Delta(G') \leq 3$ and $\text{tw}(G') \leq 2$, so $\text{cw}(G') \leq 3$ by the induction hypothesis. Let (T', w') be a carving decomposition of G' of width $\text{cw}(G')$. Let T be the tree obtained from T' by first changing the label of the leaf of T' corresponding to vertex v into p , and then adding two new vertices u and v and two new edges pu and p ; see Figure 2. Since v is the only neighbor of u in G , it is easy to see that the width of the resulting carving decomposition (T, w) of G is at most $\text{cw}(G')$, which implies that $\text{cw}(G) \leq \text{cw}(G') \leq 2$.




Fig. 2. A schematic illustration of how the tree T is constructed from the tree T' in the proof of Theorem 1.

Finally, suppose that G contains a vertex u of degree 2. Since we assume G to be simple, u has two distinct neighbors v and t . Let $G' = (V', E')$ denote the connected graph obtained from G by dissolving u . Note that G' has maximum degree at most 3, and that $\text{tw}(G') \leq 2$ due to Lemma 2 and the fact that G' is a minor of G . Hence, by the induction hypothesis, $\text{cw}(G') \leq 3$. If $\text{cw}(G') \leq 1$, then $\Delta(G') \leq 1$ by Observation 1. This, together with the observation that G' is a

connected graph on $|V| - 1 \geq 2$ vertices, implies that G' is a path on two vertices. Consequently, G is a path on three vertices, and hence $\text{cw}(G) = 2 \leq 3$. If $2 \leq \text{cw}(G') \leq 3$, then we can apply Lemma 6 to conclude that $\text{cw}(G) = \text{cw}(G') \leq 3$. This completes the proof of Theorem 1. \square

Since graphs of treewidth at most 2 can easily be recognized in linear time, Theorem 1 implies a linear-time recognition algorithm for graphs of carving-width at most 3.

Thilikos, Serna and Bodlaender [20] proved that for any k , there exists a linear-time algorithm for constructing the immersion obstruction set for graphs of carving-width at most k . For $k \in \{1, 2\}$, finding such a set is trivial. We now present an explicit description of the immersion obstruction set for graphs of carving-width at most 3.




Fig. 3. The immersion obstruction set for graphs of carving-width at most 3.

Corollary 1. *A graph has carving-width at most 3 if and only if it does not contain any of the six graphs in Figure 3 as an immersion.*

Proof. Let G be a graph. We first show that if G contains one of the graphs in Figure 3 as an immersion, then G has carving-width at least 4. In order to see this, it suffices to observe that the graphs K_4, H_1, \dots, H_5 all have carving-width 4. Hence, G has carving-width at least 4, because carving-width is a parameter that is closed under taking immersions (cf. [20]).

Now suppose that G has carving-width at least 4. Then, due to Theorem 1, $\Delta(G) \geq 4$ or $\text{tw}(G) \geq 3$. If $\Delta(G) \geq 4$, then G has a vertex v of degree at least 4. By considering v and four of its incident edges, it is clear that G contains one of the graphs H_1, \dots, H_5 as a subgraph, and consequently as an immersion. If $\text{tw}(G) \geq 3$, then Lemma 3 implies that G contains K_4 as a topological minor, and consequently as an immersion. \square

From the proof of Corollary 1, we can observe that an alternative version of Corollary 1 states that a graph has carving-width at most 3 if and only if it does not contain any of the six graphs in Figure 3 as a topological minor.

4 Conclusions

Extending Theorem 1 to higher values of carving-width remains an open problem, and finding the immersion obstruction set for graphs of carving-width at

most 4 already seems to be a challenging task. We proved that for any graph G , $\text{cw}(G) \leq 3$ if and only if $\Delta(G) \leq 3$ and $\text{tw}(G) \leq 2$. We finish our paper by showing that the equivalence “ $\text{cw}(G) \leq 4$ if and only if $\Delta(G) \leq 4$ and $\text{tw}(G) \leq 3$ ” does not hold in either direction.

To show that the forward implication is false, we consider the pentagonal prism F_5 , which is displayed in Figure 4 together with a carving decomposition (T, w) of width 4. Hence, $\text{cw}(F_5) \leq 4$. However, F_5 is a minimal obstruction for graphs of treewidth at most 3 [4], implying that $\text{tw}(F_5) = 4$.




Fig. 4. The pentagonal prism F_5 and a carving decomposition (T, w) of F_5 that has width 4.

To show that the backward implication is false, we consider the graph K_5^- , which is the graph obtained from K_5 by removing an edge. Note that $\Delta(K_5^-) = 4$ and $\text{tw}(K_5^-) = 3$. It is not hard to verify that $\text{cw}(K_5^-) = 6$. Since removing an edge decreases the carving-width by at most 1, we conclude that $\text{cw}(K_5^-) \geq 5$.

References

1. Abu-Khzam, F.N., Langston, M.A.: Graph coloring and the immersion order. In: Warnow, T., Zhu, B. (eds.) COCOON 2003. LNCS, vol. 2697, pp. 394–403. Springer (2003)
2. Arnborg, S., Proskurowski, A.: Characterization and recognition of partial 3-trees. SIAM J. Algebraic Discrete Methods 7(2), 305–314 (1986)
3. Belmonte, R., van ’t Hof, P., Kamiński, M., Paulusma, D., Thilikos, D.M.: Characterizing graphs of small carving-width. In: Lin, G. (ed.) COCOA 2012. LNCS, vol. 7402, pp. 360–370. Springer (2012)
4. Bodlaender, H.L.: A partial k -arboretum of graphs with bounded treewidth. Theoretical Computer Science 209(1-2), 1–45 (1998)
5. Bodlaender, H.L., Thilikos, D.M.: Graphs with branchwidth at most three. Journal of Algorithms 32(2), 167–194 (1999)
6. DeVos, M., Dvořák, Z., Fox, J., McDonald, J., Mohar, B., Scheide, D.: Minimum degree condition forcing complete graph immersion. CoRR, arXiv:1101.2630, January 2011.

7. Diestel, R.: Graph Theory. Springer-Verlag, Electronic Edition (2005)
8. Dvořák, Z., Giannopoulou, A.C., Thilikos, D.M.: Forbidden graphs for tree-depth. *European Journal of Combinatorics* 33(5), 969–979 (2012)
9. Giannopoulou, A.C., Kamiński, M., Thilikos, D.M.: Forbidding Kuratowski graphs as immersions. Submitted for publication.
10. Grohe, M., Kawarabayashi, K., Marx, D., Wollan, P.: Finding topological subgraphs is fixed-parameter tractable. In: *Proceedings of STOC 2011*, pp. 479–488, ACM (2011)
11. Kawarabayashi, K., Kobayashi, Y.: List-coloring graphs without subdivisions and without immersions. In: *Proceedings of SODA 2012*, pp. 1425–1435, SIAM (2012)
12. Khuller, S., Raghavachari, B., Young, N.: Designing multicommodity flow trees. *Information Processing Letters* 50, 49–55 (1994)
13. Koutsonas, A., Thilikos, D.M., Yamazaki, K.: Outerplanar obstructions for matroid pathwidth. *Electronic Notes in Discrete Mathematics* 38, 541–546 (2011)
14. Robertson, N., Seymour, P.D., Thomas, R.: Linkless embeddings of graphs in 3-space. *Bull. Amer. Math. Soc.* 28, 84–89 (1993)
15. Robertson, N., Seymour, P.D.: Graph minors XXIII. Nash-Williams’ immersion conjecture. *Journal of Combinatorial Theory, Series B* 100, 181–205 (2010)
16. Rué, J., Stavropoulos, K.S., Thilikos, D.M.: Outerplanar obstructions for a feedback vertex set. *European Journal of Combinatorics* 33, 948–968 (2012)
17. Seymour, P.D., Thomas, R.: Call routing and the ratcatcher. *Combinatorica* 14(2), 217–241 (1994)
18. Takahashi, A., Ueno, S., Kajitani, Y.: Minimal acyclic forbidden minors for the family of graphs with bounded path-width. *Discrete Mathematics* 127, 293–304 (1994)
19. Thilikos, D.M.: Algorithms and obstructions for linear-width and related search parameters. *Discrete Applied Mathematics* 105, 239–271 (2000)
20. Thilikos, D.M., Serna, M.J., Bodlaender, H.L.: Constructive linear time algorithms for small cutwidth and carving-width. In: Lee, D.T., Teng, S.-H. (eds.) *ISAAC 2000*. LNCS, vol. 1969, pp. 192–203. Springer (2000)
21. Wollan, P.: The structure of graphs not admitting a fixed immersion. Submitted for publication. Preprint available at <http://arxiv.org/abs/1302.3867>.